




Saturn Drifting Bursts (SDBs)

by
Ulrich Taubenschuss



RPWS Spectrum

RPWS: Radio and Plasma Wave Science Experiment
(3 electric monopoles,
3 magnetic search coils)

first SDB detection by Cassini
on 2004/ DOY 184 (July 2)
after SOI maneuver

named SDBs – "Saturn
Drifting Bursts"

from 3 - 60 kHz (HFR)

negative/positive drift
(negative: 88%)


Sporadic spectral occurrence

SDB storm duration: 1 – 10
hours (max. 43 hours)

fundamental emission
→ LH O-mode

1st harmonic emission
(n=2) → RH X-mode

fundamental & harmonic
visible together in ~20% of
cases




Time – Frequency Statistics

Analyzed period: 2004 (SOI) – end of 2011 4.5 years of data

11252 SDBs identified (88% negative drift; 12% positive drift)

60153 spectral points scanned – various parameters derived for scanned points (time, frequency, intensity, Cassini-location)

Fundamental & harmonic distinguishable in 20% of cases (when visible together)




most probable fundamental frequency:

- negative drift ... 9 kHz
- positive drift ... 15 kHz

highest fundamental frequencies:


- negative drift ... 20 kHz
- positive drift 25 / 30 kHz



max. temporal extension:

15 – 20 minutes

peak: 1 – 2 minutes




fundamental frequency extension:

< 1 – 7 kHz

peak: 1 – 2 kHz

Time – Frequency Statistics

Linear drift rate:



linear drift-rate (kHz/min) is a **function of frequency!**


range: 0.1 – 2 kHz/min

there is a small residual quadratic drift (kHz/min²) → curvature of SDB

quadratic drift too small to be analyzed (limited frequ. resolution of HFR receiver)


Meridional plane:

mean-frequency per bin has no special distribution with radius/latitude of Cassini



Equatorial plane:

mean-frequency per bin has no special distribution with radius/local-time of Cassini





Polarization

Direction-Finding (DF) mode: all 3 electric antennas active (U-, V-, W-antenna) ~ 17% of all data (rest = Dipole-mode data)

→ yields 3 auto-correlation signals: $\langle UU^* \rangle$, $\langle VV^* \rangle$, $\langle WW^* \rangle$

→ yields 2 real and 2 imaginary parts of cross-correlations: $\langle \text{Re}\{UW^*\} \rangle$, $\langle \text{Im}\{UW^*\} \rangle$, $\langle \text{Re}\{VW^*\} \rangle$, $\langle \text{Im}\{VW^*\} \rangle$


7 signals are enough to determine 6 unknown parameters: **Stokes parameters** (S, Q, U, V) and **direction of incidence** (θ , ϕ)
(works if **E** perp. **k** ($f_{\text{wave}} \gg f_{\text{cutoff}}$); if signal at least 20 – 30 dB above background; if **k** out ($> 10^\circ$) of antenna plane)




SDBs are strongly **circularly polarized**: d_c peaks at 0.8

high total polarization:
 d peaks at 0.85

Northern hemisphere:
- fundamental appears LH
- harmonic appears RH




Southern hemisphere:
- fundamental appears RH
- harmonic appears LH



Polarization with regard to magnetic field:

- **fundamental ... LH O-mode**
- **harmonic RH X-mode**

Statistics: Position of Cassini



- SDB visibility is strongly varying during 4.5 years (mid of 2004 – end of 2011) [SDB storms](#)
- more SDBs seen by Cassini during high inclination orbits

Latitude:

Visibility as a function of (chronographic) **latitude** of Cassini

Count number of SDBs in spectra per LAT-bin

Importance of **normalization** of visibility-histograms

Lowest visibility in the equatorial plane

2 main peaks:

-) around +40° north LAT
-) around -40° south LAT

→ preferential **beaming of SDBs into the mid-latitude sector**





2 side peaks:

-) around +65° north LAT
-) around -70° south LAT

→ propagation effect; SDBs at 55° - 60° (sinks) are blocked from the observer by high density plasma region?

→ side peaks at high latitudes (65° - 70°) are due to enhanced O-mode count rate; O-mode (fundamental) can propagate across the poles from the other side; X-mode (harmonic) is blocked (high B over poles).

SDB Visibility (2004 – 2011)



no harmonics (X-mode) in side peaks?

SDB Visibility (2004 – 2011)

Visibility with radial distance:

-) peak around 20 Rs
-) sporadic observations beyond 50 Rs (low SNR)
-) most distant obs. at 77 Rs

Visibility with Local Time (LT):

-) 2 peaks: midnight and noon
- LT is strongly coupled with LAT – special orbit geometry
-) but, peaks with LT are also visible for observations in the equatorial plane only ($|\text{Lat}| < 5^\circ$)
- variation with LT might be real

Visibility with SLS4:


-) separate observations into north and south; $|\text{Lat}| > 15^\circ$
-) clear modulation with SLS4 in both hemispheres
-) north and south have approx. the same phase ($100^\circ - 113^\circ$)

No modulation of SDB-visibility as a function of SLS4 at LT 12:00.


"clock-like" (flash-light) source?

- increased activity if a certain SLS4-longitude (-10°) is pointing at the observer
- + narrow beaming characteristic


Radial dist. of Cassini during SDB obs.




Local Time of Cassini during SDB obs.



SLS4 of Cassini during SDB obs.



SLS4 at LT 12:00 of Cassini during SDB obs.



SDB Visibility (2004 – 2011)

Visibility summary:

-) preferential beaming of radiation into **mid latitudes** ($\pm 40^\circ$)



-) variation with Local Time; mostly observed around **midnight** and **noon**

-) visibility out to large radial distances (**80 R_s**); visibility outside Saturn's magnetosphere $\rightarrow$ evidence of electro-magnetic radiation



-) modulation with **SLS4**; higher SDB activity around SLS4 = $-10^\circ/-20^\circ$ north/south $\rightarrow$ "clock-like" source?

-) sporadic events; no permanent visibility; longest continuous observation ("SDB storm") lasting ~ 43 hours (with gaps < 3 h).


All Cassini locations



Cassini locations during SDB observation (normalized)




Statistics: Intensity



Meridional plane:

mean-intensity per bin has a clear distribution with radius/latitude of Cassini

- higher intensities at small radial distance ($< 10 R_S$)
- higher intensities at higher latitudes ($> 30^\circ$)




Equatorial plane:

mean-intensity per bin has no special distribution with LT of Cassini, but with radial distance $\rightarrow$ proximity to source region.

Intensity = apparent Stokes-S [$V^2 m^{-2} Hz^{-1}$]

DP-mode: $S_{app} = \text{auto}_D / I_D^2 + \text{auto}_W / I_W^2$

DF-mode: $S_{app} = \text{auto}_U / I_U^2 + \text{auto}_V / I_V^2 + \text{auto}_W / I_W^2$



S_{app} normalized by distance² is a function of frequency!


$\rightarrow$ efficiency of the generation mechanism for SDBs is a function of frequency?

Mode Conversion

Focus on electrostatic **upper hybrid oscillations** $\rightarrow$ thermally excited at $f_{UH}^2 = f_c^2 + f_p^2$

Upper hybrid oscillations can be converted to electromagnetic modes (Z-mode, O-mode) by linear (Oya, 1971; Jones, 1976) or nonlinear mechanisms (Melrose, 1981).

Focus on **linear mode conversion**: UHR $\rightarrow$ Z-mode $\rightarrow$ O-mode



Beam/anisotropy in velocity distribution/MHD wave shifts UH oscillation onto the Z-mode branch at $f \leq f_{uh}$.

Z-mode propagates into a region where $f_z = f_p$.

Z-mode converts to O-mode
at $f \approx f_p$ (1st Ellis- window).


A gradient in the plasma density is required for this scenario.

$\nabla n / \lambda$ must be in the right range for an efficient conversion.

Mode Conversion

Linear mode conversion

- 1) UHR $\rightarrow$ Z-mode
- 2) Z-mode propagates towards increasing plasma density
- 3) Z-mode is reflected at cutoff and propagates back towards decreasing plasma density
- 4) In the layer where $f_p \approx f_z$, it can **convert to O-mode** ($f_o \approx f_p \approx f_z$)
- 5) O-mode escapes towards decreasing plasma density into "free space"; detected remotely by spacecraft.
- 6) Z-mode which didn't convert gets absorbed again (Z-mode $\rightarrow$ UHR)



Mode Conversion

Sporadic conversion event 2009/DOY 207

Cassini is at medium northern latitudes: $20^\circ - 35^\circ$

SKR is dominantly RH X-mode
 → red shades RH
 → blue shades ... LH
 Sign of Im (+ vs. -) depends on direction of incidence: source below or above antenna plane.

unpolarized UHR (white imaginary-part) at $f < 11$ kHz becomes intense, bursty and circularly polarized


the sense of apparent circular polarization stays constant → systematic behavior

Interpretation: mode conversion

UHR → Z-mode
 Z-mode → O-mode

Mode conversion is active along a broad range of latitudes

Cassini is crossing source regions for SDBs?



Mode Conversion

Where does mode conversion occur?

scan spectral points where UHR becomes polarized (2004 – 2012)

data selection:

-) $S_{app} > 10^{-14} \text{ V}^2 \text{ m}^{-2} \text{ Hz}^{-1}$

-) $d_{app} > 0.5$

histograms for location of conversion (= location of Cassini)





histograms are normalized to the time Cassini spends in a certain bin; exclude positions at $r > 15 R_S$; UHR becomes invisible; $L_{Debye} \gg L_{antenna}$

Results

- Polarized UHR seen at all LT and along a broad range of latitudes (up to 50°)

- 4 groups:

- 1) sporadic conversion
- 2) conv. in injection events
- 3) conv. close to Enceladus
- 4) conv. when UHR meets ECH






- 2 peaks in occurrence of radial distances:
 $r = 2.5 - 4.0 R_S$: Enceladus flybys, UHR between ECH
 $r = 5.0 - 9.0 R_S$: sporadic events, injection events

- 2 peaks in occurrence of frequencies:
 $\rightarrow 10 \text{ kHz vs. } 40 \text{ kHz}$
 (strongly correlated with radial dist. and lat.)

Direction Finding – Case Study

2008/344 12:00:

Cassini at high northern latitudes (40°);
close to Saturn ($6.4 R_s$)






SDBs are poorly resolved in DF-mode
($\Delta t = 32\text{sec}$, $\Delta f/f \approx 5\%$)

Scan 1 point/frequency-channel along the spectral
middle line

SNR of scanned points should be $> 10^3$

Scanned points $\rightarrow$ [direction-finding](#) (S, Q, U, V, θ , ϕ)

Direction Finding – Case Study



- SDB spectral drift is generated by a **moving source** as a function of frequency
- mode conversion scenario: **SDB-frequency = f_p in crossed layer**
- source is moving along $\sim$ const. latitude towards greater radial distance $\rightarrow f_p$ decreasing $\rightarrow$ **negative drifting SDB**
- source is moving at a high inclination to dipol field lines
- source is drifting in LT, against rotation direction of MS (= drift-direction of electrons)

Conclusions

Summary:

- Saturn Drifting Bursts (SDB) detected by Cassini/RPWS below 60 kHz
- Sporadic events (storms of several hours)
- Drift rates 0.1 - 2 kHz/min (positive, negative)
- 20% of cases: fundamental & 1st harmonic visible (harmonic at twice the fundamental frequency)
- fundamental = LH O-mode; harmonic = RH X-mode (circularly polarized)
- best visibility at mid latitudes (40° north & south) around noon and midnight
- generation mechanism: linear or non-linear mode conversion ($f_{\text{UHR}} \rightarrow f_z = f_p = f_o$); explains polarization and harmonics
- spectral drift is the consequence of a source moving through layers of decreasing/increasing plasma frequency

Open questions:

- is mode conversion working efficiently enough in Saturn's plasma torus (linear or non-linear mechanism)?
- what kind of particles are providing free energy: electrons, protons or water group ions (H^+ , OH^+)?
- what kind of particle distribution function is initiating the conversion process (direct source-crossing needed)?
- what is the particle energy range needed for explaining drift durations of minutes?